

Estimating Lorenz Curve for Iran by Using Continuous L_1 Norm Estimation

Bijan Bidabad¹

Abstract

In this paper, the L_1 norm of continuous functions and corresponding continuous estimation of regression parameters are defined. The continuous L_1 norm estimation problem of one and two parameters linear models in the continuous case are solved. We proceed to use the functional form and parameters of the probability distribution function of income to exactly determine the L_1 norm approximation of the corresponding Lorenz curve of the statistical population under consideration. Iran family budget data were used to estimate income distribution for the period of 1362-1370.

Keywords: L_1 norm, Lorenz Curve, Continuous estimation, Income distribution

1. Introduction

The skewness of income distribution is persistently exhibited for different populations and at different times. It is discussed that Pearsonian family distributions are rival functions to explain income distribution. Lorenz curve is a method to analyze the skew distributions. There is a relation between the area under the Lorenz curve and the corresponding probability distribution function of the statistical population (see, Kendall and Stuart (1977)). That is, when the probability distribution function is known, we may find the corresponding Gini coefficient as the measure of inequality.

Estimation of the Lorenz curve is confronted with some difficulties. For this estimation, we should define an appropriate functional form which can accept different curvatures (see, Bidabad and Bidabad (1989a,b)). There is another problem, that is, to create the necessary data set for estimating the corresponding parameters of the Lorenz curve, a large amount of computation on raw sample income data is inevitable. Obviously, these problems, despite their computational difficulties, make the significance of the estimated parameters poor (see, Bidabad and Bidabad (1989a,b)). To avoid this, we try to estimate the functional form of the Lorenz curve by using continuous information. In this paper, we use the probability density function of population income to estimate the Lorenz function parameters. The continuous L_1 norm smoothing method, which will be developed for estimating the regression parameters is used to solve this problem. However, we concentrate on two rival probability density functions of Pareto and log-normal. Since the former is simply integrable, there is no general problem to derive the corresponding Lorenz function, and the function is uniquely derived. But in the latter case, the log-normal density function (which has better performance for full income range) than Pareto distribution (which better fits to higher income range, (see, Cramer (1973), Singh and Maddala (1976), Salem and Mount (1974)), is not integrable and we can not determine its corresponding Lorenz function. In this regard, we should solve the problem by defining a general Lorenz curve functional form and applying the L_1 norm smoothing to estimate the corresponding parameters.

In this paper, continuous L_1 norm estimation is developed by using a similar method proposed in Bidabad (1987a,88a,89a,b) for the discrete case. Then the method is applied to the estimation of the Lorenz curve functional forms which have been proposed by Gupta (1984) and Bidabad and Bidabad (1989,92). In the end, we use our formulation to estimate Gini ratio and Kakwani length indices of inequality for the United States for the period of 1971-1990, based on the assumption that income is distributed log-normally.

2. L_1 norm of continuous functions

Generally, L_p norm of a function $f(x)$ (see, Rice and White (1964)) is defined by,

$$\|f(x)\|_p = \int_{x \in I} |f(x)|^p dx^{1/p} \quad (1)$$

Where, "I" is a closed bounded set. The L_1 norm of $f(x)$ is simply written as,

$$\|f(x)\|_1 = \int_{x \in I} |f(x)| dx \quad (2)$$

Suppose that the non-stochastic function $f(x, \beta)$ of "x", is combined with stochastic disturbance term "u" to form $y(x)$ as follows,

$$y(x) = f(x, \beta) + u \quad (3)$$

¹ (B.A., M.Sc., Ph.D., Post-Doc.) Professor of Economics and Chief Islamic Banking Advisor, Bank Melli Iran, bijan@bidabad.com, Bidabad@yahoo.com, http://www.bidabad.com



Where, β is unknown parameters vector. Rewriting u as the residual of $y(x)-f(x,\beta)$, for L_1 norm approximation of " β " we should find " β " vector such that the L_1 norm of " u " is minimum. That is,

$$\text{Min: } S = \int_{x \in I} |y(x) - f(x, \beta)| dx \quad (4)$$

3. Linear one parameter L_1 norm continuous smoothing

Redefine $f(x,\beta)$ as βx and $y(x)$ as the following linear function,

$$y(x) = \beta x + u \quad (5)$$

Where, " β " is a single (non-vector) parameter. Expression (4) reduces to:

$$\text{min: } S = \int_{x \in I} |y(x) - \beta x| dx \quad (6)$$

The discrete analog of (6) is solved by Bidabad (1987a,88a,89a,b). In these papers, we proposed applying discrete and regular derivatives to the discrete problem by using a slack variable " t " as a point to distinguish negative and positive residuals. A similar approach is used here to minimize (6). To do so in this case, certain Lipschitz conditions are imposed on the functions involved (see, Usow (1967a)). Rewrite (6) as follows,

$$\text{Min: } S = \int_{x \in I} |x| |y(x)/x - \beta| dx \quad (7)$$

For convenience, define " I " as a closed interval $[0,1]$. The procedure may be applied to other intervals with no major problem (see, Usow (1967a), Hobby and Rice (1965), Kripke and Rivlin (1965)). To minimize this function, we should first remove the absolute value sign of the expression after the integral sign. Since " x " belongs to closed interval " I ", $y(x)$ (which is a linear function of " x ") and also $y(x)/x$ are smooth and continuous. Thus, since $y(x)/x$ is uniformly increasing or decreasing function of " x ", a value of $t \in I$ can be found to have the following properties,

$$\begin{aligned} y(x)/x &< \beta & \text{if } x < t \\ y(x)/x &= \beta & \text{if } x = t \\ y(x)/x &> \beta & \text{if } x > t \end{aligned} \quad (8)$$

Value of the slack variable " t " actually is the border of negative and positive residuals. If the value of " t " were known, from (8) (middle equation), we could calculate the optimal value of " β " or inversely. But nor " t " neither " β " are known. To solve this problem, according to (8), we can rewrite (7) as two separate definite integrals with different upper and lower bounds.

$$\text{min: } S = - \int_0^t |x| (y(x)/x - \beta) dx + \int_t^1 |x| (y(x)/x - \beta) dx \quad (9)$$

Decomposition of (7) into (8) has been done by use of the slack variable " t ". Since both " β " and " t " are unknown, to solve (9), we partially differentiate it with respect to " t " and " β " variables.

$$\frac{\delta S}{\delta \beta} = \int_0^t |x| dx - \int_t^1 |x| dx = 0 \quad (10)$$

and using Liebniz' rule to differentiate the integrals with respect to their variable bounds " t ", yields,

$$\frac{\delta S}{\delta t} = -|t| \left[\frac{y(t)}{t} - \beta \right] - |t| \left[\frac{y(t)}{t} - \beta \right] = 0 \quad (11)$$

Since " x " belongs to $[0,1]$, equation (10) can be written as,

$$\int_0^t x dx - \int_t^1 x dx = 0 \quad (12)$$

or,

$$\frac{1}{2} t^2 - \frac{1}{2} + \frac{1}{2} t^2 = 0 \quad (13)$$

Which yields,

$$t = \sqrt{2}/2 \quad (14)$$

Substitute for " t " in equation (11), yields,

$$\beta = \frac{y(\sqrt{2}/2)}{\sqrt{2}/2} \quad (15)$$

Remember that $y(t)$ is function $y(x)$ evaluated at $x=t$. Value of " β " given by (15) is the optimal solution of (6). The above procedure actually is a generalization of Laplace weighted median for the continuous case.

Before applying this procedure to the Lorenz curve, let us develop the procedure for the two parameters linear model.

4. Linear two parameters L1 norm continuous smoothing

Now, we try to apply the above technique to the linear two parameters model. Rewrite (4) as,

$$\text{Min: } S = \int_0^1 |y(x) - \alpha - \beta x| dx \quad (16)$$

Where, " α " and " β " are two single (non-vector) unknown parameters and $y(x)$ and " x " are as before. According to Rice (1964c), let $f(\alpha^*, \beta^*, x)$ interpolates $y(x)$ at the set of canonical points $\{x_i; i=1,2\}$, if $y(x)$ is such that $y(x) - f(\alpha^*, \beta^*, x)$ changes sign at these x_i 's and at no other points in $[0,1]$, then $f(\alpha^*, \beta^*, x)$ is the best L_1 norm approximation to $y(x)$ (see also, Usow (1967a)). With the help of this rule, if we denote these two points to t_1 and t_2 we can rewrite (16) for $I=[0,1]$ as,

$$S = \int_0^{t_1} [y(x) - \alpha - \beta x] dx - \int_{t_1}^{t_2} [y(x) - \alpha - \beta x] dx + \int_{t_2}^1 [y(x) - \alpha - \beta x] dx \quad (17)$$

Since t_1 and t_2 are also unknowns, we should minimize S with respect to α , β , t_1 and t_2 . Taking partial derivative of (17) using Leibniz' rule with respect to these variables and equating them to zero, we will have,

$$\frac{\delta S}{\delta \alpha} = - \int_0^{t_1} dx + \int_{t_1}^{t_2} dx - \int_{t_2}^1 dx = 0 \quad (18)$$

$$\frac{\delta S}{\delta \beta} = - \int_0^{t_1} x dx + \int_{t_1}^{t_2} x dx - \int_{t_2}^1 x dx = 0 \quad (19)$$

$$\frac{\delta S}{\delta t_1} = 2[y(t_1) - \alpha - \beta t_1] = 0 \quad (20)$$

$$\frac{\delta S}{\delta t_2} = -2[y(t_2) - \alpha - \beta t_2] = 0 \quad (21)$$

Equations (18) through (21) may be solved simultaneously for α , β , t_1 and t_2 . Thus, we have the following system of equations,

$$2t_2 - 2t_1 - 1 = 0 \quad (22)$$

$$t_2^2 - t_1^2 - \frac{1}{2} = 0 \quad (23)$$

$$y(t_1) - \alpha - \beta t_1 = 0 \quad (24)$$

$$y(t_2) - \alpha - \beta t_2 = 0 \quad (25)$$

The solutions are,

$$t_1 = 1/4 \quad (26)$$

$$t_2 = 3/4 \quad (27)$$

$$\alpha = y(3/4) - (3/4)\beta = y(1/4) - (1/4)\beta \quad (28)$$

$$\beta = 2[y(3/4) - y(1/4)] \quad (29)$$

This procedure, similar to that of multiple regression model for discrete case may be expanded to include " m " unknown parameters which is not discussed here. Some computational methods for solving the different cases of m parameters model are investigated by Ptak (1958), Rice and White (1964), Rice (1964a,b,c,69,85), Usow (1967a), Lazarski (1975a,b,c,77) (see also, Hobby and Rice (1965), Kripke and Rivlin (1965), Watson (1981)). Now, let us have a look at Lorenz curve and its proposed functional forms.

5. Lorenz curve

The Lorenz curve for a random variable with probability density function $f(v)$ may be defined as the ordered pair²,

$$(P(V|V \leq v), \frac{E(V|V \leq v)}{E(V)}) \quad v \in R \quad (30)$$

Where " P " and " E " stand for probability and expected value operators. For a continuous density function $f(v)$, (30) can be written as,

$$\left(\int_{-\infty}^v f(w) dw, \frac{\int_{-\infty}^v w f(w) dw}{\int_{-\infty}^v f(w) dw} \right) \equiv (x(v), y(x(v))) \quad (31)$$

² Taguchi (1972a,b,c,73,81,83,87,88) multiplies the second element of (30) by $P(V|V \leq v)$ which is not correct; his definition of (31) is equivalent to ours.

$$\int_{-\infty}^{+\infty} wf(w)dw$$

We denote (31) by $(x(v), y(x(v)))$ where $x(v)$ and $y(x(v))$ are its elements. Therefore, "x" is a function which maps "v" to $x(v)$ and "y" is a function which maps $x(v)$ to $y(x(v))$. The function $y(x(v))$ is simply the Lorenz curve function. In recent years some functional forms for the Lorenz curve have been introduced. Among different proposed functions, we use the forms of Gupta (1984) and Bidabad and Bidabad (1989,92) which benefits from certain properties (see their articles for more explanations). Gupta (1984) proposed the functional form,

$$y = xA^{x-1} \quad A > 1 \quad (32)$$

Bidabad and Bidabad (1989,92) suggest the following functional form:

$$y = x^B A^{x-1} \quad B \geq 1, A \geq 1 \quad (33)$$

To estimate the above functions by regular estimating method, we should gather discrete data from the statistical population, and manipulate them to construct relevant x and y vectors to estimate "A" of (32) or "A" and "B" of (33). If the probability distribution of income is known, instead of gathering discrete observations, we can estimate the Lorenz curve by using the continuous L_1 norm smoothing method for continuous functions. In the following section, we proceed to apply this method to estimate the parameters "A" of (32) and "A" and "B" of (33) by using the information of probability density function of income.

6. Continuous L_1 norm smoothing of Lorenz curve

To estimate the Lorenz curve parameters when income probability density function is known, we can not always take straightforward steps. When the probability density function is easily integrable, there is no major problem in advance. We can find the functional relationship between the two elements of (31) by simple mathematical derivation. But, when integrals of (31) are not obtainable, another procedure should be adopted.

Suppose that the income of society is distributed with probability density function $f(w)$. This density function may be a skewed function such as Pareto or log-normal, as follows

$$f(w) = \theta k \theta w^{-\theta-1}, \quad w > 0, \theta > 0 \quad (34)$$

$$f(w) = [1/w\sigma\sqrt{2\pi}] \exp\{-[\ln(w)-\mu]^2/2\sigma^2\}, \quad w \in (0, \infty), \mu \in (-\infty, +\infty), \sigma > 0 \quad (35)$$

These two distributions have been known as good candidates for presenting the distribution of personal income.

In the case of Pareto density function of (34), we can simply derive the Lorenz curve function as follows. Let $F(w)$ denote the Pareto distribution function:

$$F(w) = 1 - (k/w)^\theta \quad (36)$$

with mean equal to,

$$E(w) = \theta^k / (\theta - 1), \quad \theta > 1 \quad (37)$$

If we find the function y as stated by (31) as a function of x, the Lorenz function will be derived. Now, proceed as follows. Rearrange the terms of (31) as,

$$x(v) = \int_{-\infty}^v f(w)dw \quad (38)$$

$$y(x(v)) = [1/E(x)] \int_{-\infty}^{tv} wf(w)dw \quad (39)$$

Substitute Pareto distribution function,

$$x(v) = F(v) = 1 - (k/v)^\theta \quad (40)$$

$$y(x(v)) = [(\theta-1)/\theta^k] \int_k^v \theta k^\theta w^{\theta-1} dw \quad (41)$$

or,

$$y(x(v)) = 1 - (k/v)^{\theta-1} \quad (42)$$

Now, by solving (40) for "v" and substituting in (42), the Lorenz curve for Pareto distribution is derived as,

$$y = 1 - (1-x)^{(\theta-1)/\theta} \quad (43)$$

As it was shown in the case of Pareto distribution, formula of Lorenz curve is easily obtained. But, if we select the log-normal density function (35), the procedure may not be the same. Because the integral of log-normal function has not been derived yet. In the following pages, the L_1 norm smoothing technique will be developed to estimate the parameters of given functional forms (32) and (33) by using the continuous probability density function.

According to (30) and (31) independent and dependent variables of (32) and (33) may be written as,

$$x(v) = \int_0^v f(w)dw \quad (44)$$

$$y(x(v)) = [1/E(x)] \int_0^v wf(w)dw \quad (45)$$

Substitute (44) and (45) inside (32) and define random error term u as,

$$\left[\frac{1}{E(w)} \int_0^v w f(w) dw \right] = \int_0^v f(w) dw \cdot A \cdot e^u \quad (46)$$

or briefly,

$$y(x) = x A^{x-1} e^u \quad (47)$$

Similarly for the model (35),

$$\left[\frac{1}{E(w)} \int_0^v w f(w) dw \right] = \left\{ \int_0^v f(w) dw \right\} \cdot A^B \cdot e^u \quad (48)$$

or briefly,

$$y(x) = x^B A^{x-1} e^u \quad (49)$$

Taking natural logarithm of (47) and (49), gives,

$$\ln y(x) = \ln x + (x-1) \ln A + u \quad (50)$$

$$\ln y(x) = B \ln x + (x-1) \ln A + u \quad (51)$$

With respect to properties of Lorenz curve and probability density function of $f(w)$ and equations (46) to (49), it is obvious that x belongs to the interval $[0,1]$. Thus the L_1 norm objective function for minimizing (50) or (51) is given by,

$$\min: S = \int_0^1 |u| dx \quad (52)$$

Now, let us deal with L_1 norm estimation of "A" of Lorenz curve functional form (32) (redefined by (50)). The corresponding L_1 norm objective function will be,

$$\min: S = \int_0^1 |\ln y(x) - \ln x - (x-1) \ln A| dx \quad (53)$$

or,

$$\min: S = \int_0^1 |x-1| \left| \frac{[\ln y(x) - \ln x]}{(x-1)} - \ln A \right| dx \quad (54)$$

By a similar technique used by (9), we can rewrite (54) as,

$$\min: S = \int_0^1 |x-1| \left\{ \frac{[\ln y(x) - \ln x]}{(x-1) - \ln A} \right\} dx - \int_0^1 |x-1| \left\{ \frac{[\ln y(x) - \ln x]}{(x-1) - \ln A} \right\} dx \quad (55)$$

since, $0 \leq x \leq 1$ we have,

$$\min: S = - \int_0^1 [\ln y(x) - \ln x - (x-1) \ln A] dx + \int_0^1 t [\ln y(x) - \ln x - (x-1) \ln A] dx \quad (56)$$

Differentiate (56) partially with respect to "t" and "A" and equate them to zero;

$$\frac{\delta S}{\delta t} = + \int_0^1 [(x-1)/A] dx - \int_0^1 t [(x-1)/A] dx = 0 \quad (57)$$

$$\frac{\delta S}{\delta A} = - 2 [\ln y(t) - \ln t - (t-1) \ln A] = 0 \quad (58)$$

From equation (57), we have,

$$t = 1 \pm \sqrt{2}/2 \quad (59)$$

Since "t" should belong to the interval $[0,1]$, we accept,

$$t = 1 - \sqrt{2}/2 \quad (60)$$

Substitute (60) in (58), and solve for "A", gives the L_1 norm estimation for "A" equal to,

$$A = \left[\frac{1 - \sqrt{2}/2}{y(1 - \sqrt{2}/2)} \right]^2 \quad (61)$$

Now, let us apply this procedure to another Lorenz curve functional form of (33) (redefined by (51)). Rewrite L_1 norm objective function (52) for the model (51),

$$\min_{A,B} S = \int_0^1 |\ln y(x) - B \ln x - (x-1) \ln A| dx \quad (62)$$

or,

$$\min_{A,B} S = \int_0^1 |x-1| \left| \frac{\ln y(x)}{(x-1)} - \frac{\ln x}{(x-1)} - \ln A \right| dx \quad (63)$$

The objective function (63) - by some changing on variables - is similar to (16). Thus, by a similar procedure to those of (17) through (29) we can write "S" as,

$$\begin{aligned} \min_{A,B} S = & \int_0^{t_1} |x-1| \left\{ \frac{\ln y(x)}{(x-1)} - \frac{\ln x}{(x-1)} - \ln A \right\} dx \\ & - \int_{t_1}^{t_2} |x-1| \left\{ \frac{\ln y(x)}{(x-1)} - \frac{\ln x}{(x-1)} - \ln A \right\} dx \\ & + \int_{t_2}^1 |x-1| \left\{ \frac{\ln y(x)}{(x-1)} - \frac{\ln x}{(x-1)} - \ln A \right\} dx \end{aligned} \quad (64)$$

Since $0 \leq x \leq 1$, then (64) reduces to,

$$\begin{aligned} \min_{A,B} S = & - \int_0^{t_1} [\ln y(x) - B \ln x - (x-1) \ln A] dx + \int_{t_1}^{t_2} [\ln y(x) - B \ln x - (x-1) \ln A] dx \\ & - \int_{t_2}^1 [\ln y(x) - B \ln x - (x-1) \ln A] dx \end{aligned} \quad (65)$$

Differentiate "S" partially with respect to "A", "B", t_1 and t_2 and equate them to zero,

$$\frac{\delta S}{\delta A} = - \left[\int_0^{t_1} (x-1) dx - \int_{t_1}^{t_2} (x-1) dx + \int_{t_2}^1 (x-1) dx \right] = 0 \quad (66)$$

$$\frac{\delta S}{\delta B} = \int_0^{t_1} \ln(x) dx - \int_{t_1}^{t_2} \ln(x) dx + \int_{t_2}^1 \ln(x) dx = 0 \quad (67)$$

$$\frac{\delta S}{\delta t_1} = -2 \{ \ln[y(t_1)] - B \ln(t_1) - (t_1-1) \ln(A) \} = 0 \quad (68)$$

$$\frac{\delta S}{\delta t_2} = 2 \{ \ln[y(t_2)] - B \ln(t_2) - (t_2-1) \ln(A) \} = 0 \quad (69)$$

The above system of simultaneous equations can be solved for the unknowns t_1 , t_2 , "A" and "B". Equation (66) is reduced to,

$$t_1^2 - t_2^2 - 2(t_1 - t_2) - 1/2 = 0 \quad (70)$$

Equation (67) can be written as,

$$t_1(\ln t_1 - 1) - t_2(\ln t_2 - 1) - 1/2 = 0 \quad (71)$$

Calculate t_1 from (70) as,

$$t_1 = 1 \pm \sqrt{(t_2^2 - 2t_2 + 3/2)} \quad (72)$$

Since $0 \leq t_1 \leq 1$, we accept,

$$t_1 = 1 - \sqrt{(t_2^2 - 2t_2 + 3/2)} \quad (73)$$

Substitute t_1 from (73) into (71), and rearrange the terms, gives;

$$\ln \frac{[1 - \sqrt{(t_2^2 - 2t_2 + 3/2)}]}{t_2^2} + t_2 - 3/2 + \sqrt{(t_2^2 - 2t_2 + 3/2)} = 0 \quad (74)$$

The root of equation (74) may be computed by a suitable numerical algorithm. However, it has been computed and rounded for five digits decimal point as,

$$t_2 = 0.40442 \quad (75)$$

Value of t_1 is derived by substituting t_2 into (73);

$$t_1 = 0.07549 \quad (76)$$

Values of "B" and "A" are computed from (68) and (69) using t_2 and t_1 given by (75) and (76). Thus,

$$B = \frac{(t_2-1)\ln y(t_1) - (t_1-1)\ln y(t_2)}{(t_2-1)\ln(t_1) - (t_1-1)\ln(t_2)} \quad (77)$$

or,

$$B = -0.84857\ln[y(0.07549)] + 1.31722\ln[y(0.40442)] \quad (78)$$

and,

$$A = [y(0.07549)]^{1.28986} [y(0.40442)]^{-3.68126} \quad (79)$$

Now, let us describe how equation (61) for the model (32) and equations (78) and (79) for the model (33) can be used to estimate the parameters of the Lorenz curve when the probability distribution function is known. In the model (32) we should solve (44) for $x(v)=1-\sqrt{2}/2$. On the other hand, we should find value of "v" such that,

$$x(v) = \int_0^v f(w)dw = 1-\sqrt{2}/2 \quad (80)$$

By substituting this value of "v" into (45), value of $y(1-\sqrt{2}/2)$ is computed. The value $y(1-\sqrt{2}/2)$ is used to compute the parameter "A" given by (61) for model (32).

The procedure for the model (33) is also similar, with the difference that two values of "v" should be computed. Once two different values of "v" are computed as follow,

$$x(v) = \int_0^v f(w)dw = 0.07549 \quad (81)$$

$$x(v) = \int_0^v f(w)dw = 0.40442 \quad (82)$$

Values of "v" are substituted in (45) to find $y(0.07549)$ and $y(0.40442)$. These values of "y" are used to compute the parameters of the model (33) by substituting them into (78) and (79).

The only problem remains is computation of related definite integrals of $x(v)$ defined by (80), (81) and (82) which can be done by appropriate numerical methods such as the enclosed sample computer program coded for MathCAD II for a complete example.

7. Income distribution in Iran

In order to compute the Lorenz curve for Iran, we try to apply the above procedure for both (32) and (33) propositions and using log-normal distribution function assumption. The source of data is "Statistical Center of Iran" who computed the mean and variance of income for urban and rural families for the period of 1362-1370 (1983-1991) from "Family Budget Surveys" of different years. These data are given in Table I. The amounts of mean and variance of income were used to derive the log-normal density function parameters μ and σ . The explained procedure of estimation then applied to the series of data of table I, and corresponding results are reported in Table 2. A sample computer program is also enclosed at the end of these pages.

Table I.

Year	Urban Data			Rural Data		
	Sample size	Income Mean	Income variance	Sample size	Income Mean	Income variance
1362	14747	918423	1106199100048	12440	471942	192638591017
1363	14728	1034169	1174389430497	12420	524623	351371674839
1364	13976	1037084	1792475461430	13587	531098	301917047049
1365	2745	1126638	1300389710415	3015	568557	404222563256
1366	2748	1147497	1410976253551	3018	710145	491696298459
1367	3987	1360121	2551576757245	4331	908530	1743056317121
1368	5492	1505970	4786980002705	6028	1052371	1019597224716
1369	9095	2010471	12587903327408	9348	1251060	5529127350603
1370	9168	2840790	66958717265779	9504	1563116	7505679968729

Source: Statistical Center of Iran.

Table 2.

Year	Gupta Model			Bidabad Model			
	A	Gini	Kakwani	A	B	Gini	Kakwani
Urban estimation							
1362	7.259	0.430	0.163	5.314	1.211	0.426	0.161
1363	6.279	0.409	0.148	4.620	1.204	0.405	0.146
1364	8.915	0.457	0.183	6.500	1.217	0.453	0.181
1365	5.943	0.401	0.143	4.385	1.202	0.397	0.141
1366	6.158	0.407	0.146	4.535	1.203	0.402	0.144
1367	7.574	0.436	0.167	5.539	1.212	0.432	0.165
1368	11.021	0.482	0.203	8.034	1.223	0.480	0.202
1369	15.841	0.522	0.236	11.676	1.227	0.521	0.236
Rural estimation							
1370	42.211	0.605	0.313	33.118	1.261	0.607	0.316
1362	5.220	0.382	0.129	3.878	1.195	0.377	0.127
1363	7.099	0.427	0.160	5.201	1.210	0.423	0.159
1364	6.152	0.406	0.146	4.531	1.203	0.402	0.144
1365	6.978	0.424	0.159	5.115	1.209	0.420	0.157
1366	5.718	0.396	0.139	4.227	1.200	0.391	0.137
1367	11.025	0.482	0.203	8.037	1.223	0.480	0.202
1368	5.472	0.389	0.134	4.054	1.198	0.384	0.132
1369	17.955	0.534	0.247	13.258	1.227	0.533	0.247
1370	15.683	0.521	0.235	11.518	1.227	0.519	0.234

References

- Bidabad, Bijan (1987a) Least absolute error estimation. The First International Conference on Statistical Data Analysis Based on the L_1 norm and Related Methods, Neuchatel, Switzerland. <http://www.bidabad.com/doc/lae-I.pdf>
- Bidabad, Bijan (1987b) Least absolute error estimation, part II. Submitted to the First International Conference on Statistical Data Analysis Based on the L_1 norm and Related Methods, Neuchatel, Switzerland. <http://www.bidabad.com/doc/lae-II.pdf>
- Bidabad, Bijan (1988a) A proposed algorithm for least absolute error estimation. Proc. of the Third Seminar of Mathematical Analysis. Shiraz Univ., 24-34, Shiraz, Iran.
- Bidabad, Bijan (1988b) A proposed algorithm for least absolute error estimation, part II. Proc. of the Third Seminar of Mathematical Analysis, Shiraz Univ., 35-50, Shiraz, Iran.
- Bidabad, Bijan (1989a) Discrete and continuous L_1 norm regressions, proposition of discrete approximation algorithms and continuous smoothing of concentration surface, Ph.D. thesis, Islamic Azad Univ., Tehran, Iran. <http://www.bidabad.com/doc/L1-norm-thesis-en.pdf>
- Bidabad, Bijan (1989b) Discrete and continuous L_1 norm regressions, proposition of discrete approximation algorithms and continuous smoothing of concentration surface, Ph.D. thesis, Islamic Azad Univ., Tehran, Iran. Farsi translation. <http://www.bidabad.com/doc/L1-norm-thesis-fa.pdf>
- Bidabad, Bijan (2005). L_1 norm based computational algorithms. <http://www.bidabad.com/doc/II-article6.pdf>
- Bidabad, Bijan (2005). L_1 norm solution of overdetermined system of linear equations. <http://www.bidabad.com/doc/II->

article5.pdf

Bidabad, Bijan (2005). L_1 norm based data analysis and related methods. <http://www.bidabad.com/doc/11-articl1.pdf>

Bidabad, Bijan (2005). New algorithms for the L_1 norm regression. <http://www.bidabad.com/doc/11-article2.pdf>

Bidabad, Bijan (2005). Comparative study of the L_1 norm regression algorithms. <http://www.bidabad.com/doc/11-articl3.pdf>

Bidabad, Bijan (2005). Continuous L_1 norm estimation of Lorenz curve. <http://www.bidabad.com/doc/11-articl4.pdf>

Bidabad, Bijan (2005). Continuous L_1 norm estimation of Lorenz curve when probability density function is known.

Bidabad, Bijan (2005). USA Income distribution counter-business-cyclical trend (Estimating Lorenz curve using continuous L_1 norm estimation). First meeting of the Society for the Study of Economic Inequality (ECINEQ), Palma de Mallorca, Spain, July 20-22, 2005.

<http://www.uib.es/congres/ecopub/ecineq/general.html>

<http://www.uib.es/congres/ecopub/ecineq/papers/039Bidabab.pdf>

<http://www.bidabad.com/doc/estimating-lorenz-us.pdf>

Bidabad, Bijan, Hamid Shahrestani. (2008) An implied inequality index using L_1 norm estimation of Lorenz curve. Global Conference on Business and Finance Proceedings. Mercedes Jalbert, managing editor, ISSN 1931-0285 CD, ISSN 1941-9589 Online, Volume 3, Number 2, 2008, The Institute for Business and Finance Research, Ramada Plaza Herradura, San Jose, Costa Rica, May 28-31, 2008, pp. 148-163. Global Journal of Business Research, Vol. 4, No. 1, 2010, pp.29-45.

<http://www.bidabad.com/doc/LI-Implied-inequality-index-4.pdf>

<http://www.bidabad.com/doc/SSRN-id1631861.pdf>

Bidabad, Bijan, Functional form for estimating the Lorenz curve, Australasian Econometric meeting, Australian National University, Australia, 1989. American Finance & Banking Review, 4(1), 17-21, 2019.

<https://www.cribfb.com/journal/index.php/amfbr/article/view/286>

<http://www.bidabad.com/doc/functional-form-lorenz.pdf>

<http://www.bidabad.com/doc/functional-form-lorenz.pptx>

Bidabad, Bijan, USA Income distribution counter-business-cyclical trend (Estimating Lorenz curve using continuous L_1 norm estimation). First meeting of the Society for the Study of Economic Inequality (ECINEQ), Palma de Mallorca, Spain, July 20-22, 2005.

<http://www.bidabad.com/doc/estimating-lorenz-us.pdf>

Cramer J.S. (1973) Empirical econometrics. North-Holland, Amsterdam.

Gupta M.R. (1984) Functional forms for estimating the Lorenz curve. *Econometrica*, 52, 1313-1314.

Hobby C.R., J.R. Rice (1965) A moment problem in L_1 approximation. *Proc. Amer. Math. Soc.*, 16, 665-670.

Kakwani N.C. (1980) Income inequality and poverty. New York, Oxford University Press.

Kakwani N.C. (1980) Functional forms for estimating the Lorenz curve: a reply. *Econometrica*, 48, 1063-64.

Kakwani N.C., N. Podder (1976) Efficient estimation of the Lorenz curve and associated inequality measures from grouped observations. *Econometrica* 44, 137-148.

- Kendall M., A. Stuart (1977) The advanced theory of statistics. vol.I, Charles Griffin & Co., London.
- Kripke B.R., T.J. Rivlin (1965) Approximation in the metric of $L_1(X,u)$. Trans. Amer. Math. Soc., 119, 101-22.
- Lazarski E. (1975a) Approximation of continuous functions in the space L_1 . Automatika, 487, 85-93.
- Lazarski E. (1975b) The approximation of the continuous function by the polynomials of power functions in L_1 space. Automatika, 487, 95-106.
- Lazarski E. (1975c) On the necessary conditions of the uniqueness of approximation by the polynomials of power functions in L_1 space. Automatika, 487, 107-117.
- Lazarski E. (1977) Approximation of continuous functions by exponential polynomials in the L_1 space. Automatika, 598, 82-87.
- Ptak V. (1958) On approximation of continuous functions in the metric $\int_a^b |x(t)| dt$ Czechoslovak Math. J. 8(83), 267-273.
- Rasche R.H., J. Gaffney, A.Y.C. Koo, N. Obst (1980) Functional forms for estimating the Lorenz curve. Econometrica, 48, 1061-1062.
- Rice J.R. (1964a) On computation of L_1 approximations by exponentials, rationals, and other functions. Math. Comp., 18, 390-396.
- Rice J.R. (1964b) On nonlinear L_1 approximation. Arch. Rational Mech. Anal., 17 61-66.
- Rice J.R. (1964c) The approximation of functions, vol. I, linear theory. Reading Mass., Addison-Wesley.
- Rice J.R. (1969) The approximation of functions, vol. II, linear theory. Reading Mass., Addison-Wesley.
- Rice J.R. (1985) Numerical methods, software, and analysis. McGraw- Hill, ch. 11.
- Rice J.R., J.S. White (1964) Norms for smoothing and estimation. SIAM Rev., 6, 243-256.
- Salem A.B.Z., T.D. Mount (1974) A convenient descriptive model of income distribution: the gamma density. Econometrica, 42, 1115-1127.
- Singh S.K., G.S. Maddala (1976) A function for the size distribution of income. Econometrica, 44, 963-970.
- Slottje D.J. (1989) The structure of earnings and the measurement of income inequality in the U.S., North-Holland Publishing Company, Amsterdam.
- Taguchi T. (1972a) On the two-dimensional concentration surface and extensions of concentration coefficient and Pareto distribution to the two dimensional case-I. Annals of the Inst. of Stat. Math., vol. 24, no.2, 355-381.
- Taguchi T. (1972b) On the two-dimensional concentration surface and extensions of concentration coefficient and Pareto distribution to the two dimensional case-II. Annals of the Inst. of Stat. Math., vol. 24, no.3, 599-619.
- Taguchi T. (1972c) Concentration polyhedron, two dimensional concentration coefficient for discrete type distribution and some new correlation coefficients etc. The Inst. of Stat. Math., 77-115.
- Taguchi T. (1973) On the two-dimensional concentration surface and extensions of concentration coefficient and Pareto distribution to the two dimensional case-III. Annals of the Inst. of Stat. Math., vol. 25, no.1, 215-237.
- Taguchi T. (1974) On Fechner's thesis and statistics with norm p. Ann. of the Inst. of Stat. Math., vol. 26, no.2, 175-193.
- Taguchi T. (1978) On a generalization of Gaussian distribution. Ann. of the Inst. of Stat. Math., vol. 30, no.2, A, 211-242.
- Taguchi T. (1981) On a multiple Gini's coefficient and some concentrative regressions. Metron, vol. XXXIX - N.1-2, 5-98.

- Taguchi T. (1983) Concentration analysis of bivariate Paretoan distribution. Proc. of the Inst. of Stat. Math., vol. 31, no.1, 1-32.
- Taguchi T. (1987) On the structure of multivariate concentration. Submitted to the First International Conference on Statistical Data Analysis Based on the L_1 Norm and Related Methods, Neuchatel, Switzerland.
- Taguchi T. (1988) On the structure of multivariate concentration - some relationships among the concentration surface and two variate mean difference and regressions. CSDA, 6, 307-334.
- Usow K.H. (1967a) On L_1 approximation: computation for continuous functions and continuous dependence. SIAM J. of Numer. Anal., 4, 70-88.
- Watson G.A. (1981) An algorithm for linear L_1 approximation of continuous functions. IMA J. Num. Anal., 1, 157-167.

Copyrights

Copyright for this article is retained by the author(s), with first publication rights granted to the journal. This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).